

Mahan, H. D. 1956. Nocturnal predation on Song Sparrow eggs by milksnake. *Wilson Bulletin* 68: 245.

Mitchell, J. C. 1994. *The Reptiles of Virginia*. Smithsonian Institution Press, Washington, DC. 352 pp.

Palmer, W. M., & A. L. Braswell. 1995. *Reptiles of North Carolina*. University of North Carolina Press, Chapel Hill, NC. 412 pp.

Stejneger, L., & T. B. Barbour. 1917. A checklist of North American amphibians and reptiles. Harvard University Press, Harvard, MA. 125 pp.

Surface, H. A. 1906. The serpents of Pennsylvania. *Pennsylvania State Department of Agriculture Monthly Bulletin, Division of Zoology* 4: 113-202.

Tennant, A., & R. D. Bartlett. 2000. *Snakes of North America, Eastern and Central Regions*. Gulf Publishing Company, Houston, TX. 587 pp.

Tryon, B. W. 1982. Additional instances of multiple egg-clutch production in snakes. *Transactions of the Kansas Academy of Science* 87: 98-104.

Uhler, F. M., C. Cottam, & T. E. Clarke. 1939. Food of snakes of the George Washington National Forest, Virginia. *Transactions of the Fourth North American Wildlife Conference* 1939: 605-622.

Williams, K. L. 1988. Systematics and Natural History of the American Milk Snakes, *Lampropeltis triangulum*. Milwaukee Public Museum, Milwaukee, WI. 176 pp.

Joseph C. Mitchell  
Department of Biology  
University of Richmond  
Richmond, Virginia 23173

Carrie Seltzer<sup>1</sup>  
University of Virginia's Blandy Experimental Farm  
The State Arboretum of Virginia  
400 Blandy Farm Lane  
Boyce, Virginia 22620

<sup>1</sup>Present address:

Carrie Seltzer  
4302 12th Place NE  
Washington, DC 20017

*Banisteria*, Number 28, 2006

© 2006 by the Virginia Natural History Society

OPPORTUNISTIC SCAVENGING BY EASTERN AMERICAN TOAD (*BUFO AMERICANUS AMERICANUS*) TADPOLES ON A DEAD RED-SPOTTED NEWT (*NOTOPHTHALMUS VIRIDESCENS VIRIDESCENS*) – The diet of most tadpoles is assumed by many to consist of algae, detritus, and protozoa, but many tadpole species supplement this herbaceous diet with animal protein (Alford, 1999). On 14 April 2005 at 1800 h EDT, one of us (JDG) observed a feeding aggregation of Eastern American Toad (*Bufo americanus americanus*) tadpoles consuming a dead Red-spotted Newt (*Notophthalmus viridescens viridescens*). The shallow pond, located in White Oak Mountain Wildlife Management Area, Pittsylvania County, Virginia (36° 46' 43.1" N, 79° 19' 56.4" W, NAD 83), measured 17 m by 61 m. Water temperature was 25° C. The tadpoles appeared to be concentrated around two openings in the body wall of the newt. The two openings were located inferior to each forelimb. Abdominal organs were protruding out of one opening. Total length of this male newt was 84 mm. The death of the newt was likely the result of an aborted predation event. When examined, the newt showed no signs of ill health except for the two holes in its body. The newt was checked again four days later. All of its skin was completely stripped to the muscle. No tadpoles were observed feeding on the carcass at this time, although thousands of tadpoles were present in the pond. We could not find the newt the next day when the pond was last checked.



Fig. 1. American Toad tadpoles scavenging an adult male Red-spotted Newt in Pittsylvania County, Virginia.

Newts are known to have tetrodotoxin and other chemicals concentrated in granular glands in the skin (Brodie & Johnson, 1974; Duellman & Trueb, 1994). Tetrodotoxin is a potent neurotoxin and, along with other skin secretions, constitute a powerful emetic. Adult newts have one-tenth the toxicity of the juvenile eft stage (Brodie, 1968). This makes them less toxic but still unpalatable to many predators. Why were the American Toad tadpoles able to eat the skin of the dead adult Red-spotted Newt and not apparently be affected by the tetrodotoxin? Tadpole behavior was normal; no abnormal movements or swimming behavior were observed. *Bufo americanus* tadpoles themselves are toxic to predators at metamorphic stages of development, when they contain bufotoxin, but not during intermediate stages of growth (Brodie et al., 1978; Formanowicz & Brodie, 1982; Brodie & Formanowicz, 1987). Elucidation of the ability of *B. americanus* tadpoles to consume newt skin without apparent ill effects may reveal new physiological or pharmacological properties in these aquatic vertebrates. Tetrodotoxin may also simply decompose quickly after death allowing the tadpoles to freely consume the skin.

#### ACKNOWLEDGMENTS

We thank Dr. Susan C. Walls and two anonymous reviewers for comments on the manuscript.

#### LITERATURE CITED

- Alford, R. A. 1999. Ecology, resource use, competition, and predation. Pp. 240-278 *In* McDiarmid, R. W., & R. Altig (eds.), *Tadpoles: The Biology of Anuran Larvae*. University of Chicago Press, Chicago, IL.
- Brodie, E. D., Jr. 1968. Investigations on the skin toxin of the Red-spotted Newt, *Notophthalmus viridescens viridescens*. *American Midland Naturalist* 80: 276-280.
- Brodie, E. D., Jr., & D. R. Formanowicz, Jr. 1987. Antipredator mechanisms of larval anurans: protection of palatable individuals. *Herpetologica* 43: 369-373.
- Brodie, E. D., Jr., D. R. Formanowicz, Jr., & E. D. Brodie, III. 1978. The development of noxiousness of *Bufo americanus* tadpoles to aquatic insect predators. *Herpetologica* 34: 302-306.
- Brodie, E. D., Jr., & J. A. Johnson. 1974. Toxicity of the urodele amphibians *Taricha*, *Notophthalmus*, *Cynops*, and *Paramesotriton* (Salamandridae). *Copeia* 1974: 506-511.
- Duellman, W. E., & L. Trueb. 1994. *Biology of Amphibians*. Johns Hopkins University Press, Baltimore, MD. 696 pp.
- Formanowicz, D. R., Jr., & E. D. Brodie, Jr. 1982. Relative palatability of members of a larval amphibian community. *Copeia* 1982: 91-97.
- Jason D. Gibson  
Galileo Magnet High School  
Danville, Virginia 24541
- Joseph C. Mitchell  
Department of Biology  
University of Richmond  
Richmond, Virginia 23173